

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Childhood Blood Lead Levels and Symptoms of Attention Deficit Hyperactivity Disorder (ADHD): A Cross-Sectional Study of Mexican Children

Siying Huang, Howard Hu, Brisa N Sánchez, Karen E. Peterson, Adrienne S. Ettinger, Héctor Lamadrid-Figueroa, Lourdes Schnaas, Adriana Mercado-García, Robert O. Wright, Niladri Basu, David E. Cantonwine, Mauricio Hernández-Avila, and Martha María Téllez-Rojo

Table of Contents

Table S1. Adjusted associations between a 1- $\mu\text{g}/\text{dL}$ increase in blood lead and CRS-R outcomes in children age 6 to 13 years using segmented regression with different initial values for iterations

Table S2. Adjusted associations between a 1- $\mu\text{g}/\text{dL}$ increase in blood lead and CRS-R outcomes in children age 6 to 13 years from piecewise linear regressions using 5 $\mu\text{g}/\text{dL}$ as the breakpoint in imputed dataset (N=578)

Table S3. Adjusted associations between a 1- $\mu\text{g}/\text{dL}$ increase in blood lead and CRS-R outcomes in children age 6 to 13 years from piecewise linear regressions or simple linear regressions using complete-case data (N=362)

Table S4. Adjusted associations between a 1- $\mu\text{g}/\text{dL}$ increase in blood lead and CRS-R outcomes in children age 6 to 13 years from piecewise linear regressions with data-driven breakpoints in the complete-case data (N=361)

Table S5. Adjusted associations between a 1- $\mu\text{g}/\text{dL}$ increase in blood lead and CRS-R outcomes in children age 6 to 13 years from piecewise linear regressions using 5 $\mu\text{g}/\text{dL}$ as the breakpoint in complete-case data (N=362)

Table S6. Adjusted associations between a 1- $\mu\text{g}/\text{dL}$ increase in blood lead and CRS-R outcomes in children age 6 to 13 years from piecewise linear regressions or simple linear regressions using the samples measured from University of Michigan department of Environmental Health Sciences labs only (N=342)

Figure S1. Crude associations between a 1- $\mu\text{g}/\text{dL}$ increase in blood lead and CRS-R outcomes in children aged 6 to 13 years with locally weighted scatterplot smoothing (LOWESS) method. The X-axis represents children's blood lead levels ($\mu\text{g}/\text{dL}$) and is converted into logarithmic scale; the Y-axis represents the Conners Rating Scales-Revised (CRS-R) scores.